

[Metni yazın]

Schwarz problem for higher-order equations in a polydisc

A. Okay Celebi

Department of Mathematics, Yeditepe University,

Istanbul, Turkey

acelebi@yeditepe.edu.tr

Abstract: In this presentation, we discuss the Schwarz boundary value problem for higher order linear complex differential equations in the unit polydisc. Firstly we state the results obtained in \mathbb{C} (see for example [1]). Secondly the integral representation for functions in $\mathbb{C}^n[2,3,4]$ is improved. Then we give the solution of the model equation with homogeneous Schwarz conditions posed in a polydisc, which enables us to define an integral operator. Thus we can convert the linear complex differential equations into an integral equation. The solution is obtained via Fredholm theory. .

Keywords: Schwarz problem, Polydisc,

P.S: This is a joint work with Umit Aksoy; Atilim University, Department of Mathematics, Ankara, Turkey

References:

- [1] U. Aksoy, A. O. Celebi; A survey on the boundary value problems for complex partial differential equations, Adv. Dyn. Syst. Appl., 5(2010), 133-158.
- [2] Begehr, H. and Dzhuraev, A., An Introduction to several complex variables and partial differential equations, Pitman Monographs and Surveys1 in Pure and Applied Mathematics, Addison Wesley Longman Limited (1997).
- [3] Begehr, H., Boundary value problems in \mathbb{C} and \mathbb{C}^n , Acta Mathematica Vietnamica, 22(1997), 407-425.
- [4] Begehr, H., Dai, D.-Q. and Li, X., Integral representation formulas in polydomains, Comp. Var., 47(2002), 463-484.

\mathbb{C}^n [

\mathbb{C}^n